

Table 18. Harvest and Feeding Intervals for Soybean Herbicides

Soybean Herbicides	Days to Harvest	
	Grain	Forage
Assure II/Targa	80	Do not feed
Basagran	30	30
Basagran + 2,4-DB	60	60
Basagran + Reflex	Apply prior to bloom	Do not feed
Basagran + Cobra	90	Do not feed
Cadet	60	Do not feed
Classic	Apply 60 days before maturity	Do not feed
Clethodim	60	Do not feed
Cobra	45	Do not feed
Engenia	Apply through June 30	7
Enlist One/Duo	Apply no later than R2 stage	Do not feed
Extreme/Thunder Master	Apply prior to bloom and 85 days before harvest	Do not feed
FirstRate	70	25
Fomesafen	45	Do not feed
Fusilade DX	Apply prior to bloom	Do not feed
Glufosinate	70	Do not feed
InterMoc	Apply prior to bloom	Do not feed
Poast	75	Do not feed ¹
Perpetuo	60	Do not feed
Prefix/Vise	90	Do not feed
Pursuit	85	Do not feed
Raptor	85 and apply prior to bloom	Do not feed
Resource	60	Do not feed
Storm	50	Do not feed
Synchrony XP	Apply 60 days before maturity	Do not feed
Tavium	Apply through V4 or June 30	Do not feed (POST)
Thifensulfuron	60	Do not feed
Torment	85	Do not feed
Ultra Blazer/Avalanche Ultra	50	Do not feed
Warrant Ultra	45	Do not feed
XtendiMax	Apply up to R1 or through June 30	7

¹Soybean hay may be fed.