

Motor Vehicle Crash Deaths

How is the US doing?

On this Page

- Overview
- Problem
- Infographics
- What Can Be Done
- Issue Details

Overview

Reducing motor vehicle crash deaths was one of the great public health achievements of the 20th century for the US. However, more than 32,000 people are killed and 2 million are injured each year from motor vehicle crashes. In 2013, the US crash death rate was more than twice the average of other high-income countries. In the US, front seat belt use was lower than in most other comparison countries. One in 3 crash deaths in the US involved drunk driving, and almost 1 in 3 involved speeding. Lower death rates in other high-income countries and a high percentage of risk factors in the US suggest that we can make more progress in reducing crash deaths.

Drivers and passengers can:

- Use a seat belt in every seat, on every trip, no matter how short.
- Make sure children are always properly buckled in the back seat in a car seat, booster seat, or seat belt, whichever is appropriate for their age, height, and weight.
- Choose not to drive while impaired by alcohol or drugs, and help others do the same.
- Obey speed limits.
- Drive without distractions (such as using a cell phone or texting).

∧ Top of Page



About **90** people die each day in the US from crashes—resulting in the highest death rate among comparison countries.*

US crash deaths fell **31%** compared to an average 56% in 19 other high—income countries* from 2000-2013.

Over **18,000** lives could be saved each year if US crash deaths equaled the average rate of 19 other high—income countries.*

Problem

Motor vehicle crash deaths in the US are still too high.

There were more than **32,000** crash deaths in the US in 2013. These deaths cost more than \$380 million in direct medical costs.

Major risk factors for crash deaths in the US.

- Not using seat belts, car seats, and booster seats contributed to over 9,500 crash deaths.
- Drunk driving contributed to more than 10.000 crash deaths.

^{*}Australia, Austria, Belgium, Canada, Denmark, Finland, France, Germany, Ireland, Israel, Japan, Netherlands, New Zealand, Norway, Slovenia, Spain, Sweden, Switzerland, and the United Kingdom.

• Speeding contributed to more than 9,500 crash deaths.

Reducing major risk factors could save thousands of lives and hundreds of millions of dollars in direct medical costs each year.

Seat belts saved over 12,500 lives in the US in 2013, yet:


- The US had lower-than-average front and back seat belt use compared with other high-income countries.
- About half of drivers or passengers who died in crashes in the US weren't buckled up.

Some proven measures of best performing high-income countries.

Even when considering population size, miles traveled, and number of registered vehicles, the US consistently ranked poorly relative to other high-income countries for crash deaths. Some of the best performing countries:

- Have policies in line with best practices, including those that address:
 - Primary enforcement of seat belt laws that cover everyone in every seat.
 - Police officers can stop a vehicle and write a ticket for anyone not buckled up.
 - Requirements for car seats and booster seats for child passengers through at least age 8.
 - Blood Alcohol Concentration (BAC) levels.
 - US, Canada and the United Kingdom define drunk driving as BAC levels at 0.08% or above; all other comparison countries use lower BAC levels (0.02-0.05%).
- Use advanced engineering and technology, such as:
 - o Ignition interlocks for all people convicted of drunk driving.
 - This device keeps the vehicle from starting unless the driver has a BAC below a pre-set low limit.
 - Automated enforcement, for example, speed and red light cameras.
 - Improvements in vehicle safety and transportation infrastructure.
- Implement proven measures, such as:
 - More use of publicized sobriety checkpoints.
 - Maintain and enforce the minimum legal drinking age.

in the US and selected high-income countries



(infographic.html#graphic)

View large image and text description (infographic.html#graphic)

∧ Top of Page

Infographics

nign-income countries. Motor vehicle crash deaths in 10 comparison high-income countries, 2013 Countries with the highest and lowest reductions in crash deaths, 2000-2013 **United States** New Zealand 5.6 Spain (highest) Canada 5.4 France Japan **AVERAGE** of 19 high-Germany income countries \checkmark 56% Spain **United States (lowest) Switzerland ⇔ 31% United Kingdom** Sweden Deaths per 100,000 people SOURCE: International Road Traffic and Accident Database (IRTAD) Road Safety

Annual Report, 2015.

High-income countries with the highest percentage of crash deaths involving alcohol or speed **Alcohol** Speed Canada **Finland** 34% 42% **UNITED STATES** Denmark 31% 40% **New Zealand** Slovenia 31% 39% Australia Germany 30% 35% Slovenia Australia 30% 33% France New Zealand 29% 33% Belguim **Netherlands** 25% 30% **Finland UNITED STATES** 22% 29% Sweden 28% Austria 19% **Netherlands** 19% 26% Switzerland Note: 19 countries reported the percentage of deaths involving alcohol and 15 countries reported the percentage of deaths related to speeding. SOURCES: WHO Global Status Report on Road Safety, 2015. IRTAD Road Safety Annual Report, 2015.

(infographic.html#infographic)

Deaths per 100,000 people

SOURCE: WHO Global Status Report on Road Safety, 2015.

View larger image and text description (infographic.html#infographic)

<u>∧ Top of Page</u>

The Federal government is

- Tracking the nation's progress in reducing crash injuries and deaths. www.cdc.gov/psr/national-summary/mvi.html (http://www.cdc.gov/psr/national-summary/mvi.html)
- Evaluating and encouraging the use of proven programs and policies.
 www.thecommunityguide.org/mvoi/index.html (http://www.thecommunityguide.org/mvoi/index.html)
 www.nhtsa.gov/staticfiles/nti/pdf/812239_Countermeasures_8thEd_TT.pdf
 (http://www.nhtsa.gov/staticfiles/nti/pdf/812239_Countermeasures_8thEd_TT.pdf)
- Collaborating with and providing guidance, resources, and tools for motor vehicle injury prevention to state, local, tribal, and federal partners.
 www.cdc.gov/motorvehiclesafety/ (http://www.cdc.gov/motorvehiclesafety/)

Drivers and passengers can

- Use a seat belt in every seat, on every trip, no matter how short.
- Make sure children are always properly buckled in the back seat in a car seat, booster seat, or seat belt, whichever is appropriate for their age, height, and weight.
- Choose not to drive while impaired by alcohol or drugs, and help others do the same.
- Obey speed limits.
- Drive without distractions (such as using a cell phone or texting).

States can

- Increase seat belt use with primary enforcement seat belt laws that cover everyone in the vehicle. www.cdc.gov/psr/national-summary/mvi.html (http://www.cdc.gov/psr/national-summary/mvi.html)
- Consider requiring car seats and booster seats for children through at least age 8 or until seat belts fit
 properly.
 - www.cdc.gov/motorvehiclesafety/seatbelts/states.html (http://www.cdc.gov/motorvehiclesafety/seatbelts/states.html)
- Reduce drunk driving by:
 - Expanding publicized sobriety checkpoints.
 www.thecommunityguide.org/mvoi/AID/sobrietyckpts.html
 (http://www.thecommunityguide.org/mvoi/AID/sobrietyckpts.html)
 - Enforcing existing 0.08% blood alcohol concentration (BAC) laws and minimum legal drinking age laws.
 - www.thecommunityguide.org/mvoi/AID/mlda-laws.html (http://www.thecommunityguide.org/mvoi/AID/mlda-laws.html)
 - Considering ignition interlock requirements for people convicted of drinking and driving, starting with their first offense.
 - www.cdc.gov/motorvehiclesafety/impaired_driving/ignition_interlock_states.html (http://www.cdc.gov/motorvehiclesafety/impaired_driving/ignition_interlock_states.html)
- Support traffic safety laws with media campaigns and visible police presence to increase restraint use and

- decrease impaired driving and speeding.
- Consult CDC's Motor Vehicle Prioritizing Interventions and Cost Calculator for States (MV PICCS) to find
 out the cost of implementing up to 14 interventions and the number of injuries prevented and lives saved.
 www.cdc.gov/motorvehiclesafety/calculator/index.html
 (http://www.cdc.gov/motorvehiclesafety/calculator/index.html)
- Consider implementing Vision Zero—a road safety approach that considers death and severe injuries on our roads unacceptable and preventable.
 www.visionzeroinitiative.com (http://www.visionzeroinitiative.com)

Health care providers can

- Give patients resources about motor vehicle safety and remind them how important it is to use a seat belt on every trip.
- Counsel parents and caregivers on using age and size-appropriate car seats, booster seats, and seat belts on every trip.
 - www.cdc.gov/Motorvehiclesafety/Child_Passenger_Safety/index.html (http://www.cdc.gov/Motorvehiclesafety/Child_Passenger_Safety/index.html)
- Talk to patients about the dangers of impaired driving and help identify counseling options as needed.
- Provide parents and caregivers of teens with resources on safe teen driving. www.cdc.gov/parentsarethekey/peds/index.html
 (http://www.cdc.gov/parentsarethekey/peds/index.html)

∧ Top of Page

Issue Details

Related Pages Other Sites Science Behind the Issue

- <u>Vital Signs Issue details: Motor Vehicle Injury Prevention, Morbidity and Mortality Weekly Report (MMWR)</u>
 (http://www.cdc.gov/mmwr/volumes/65/wr/mm6526e1.htm?scid=mm6526e1 w)
- <u>Vital Signs Motor Vehicle Crash Deaths [PODCAST 1:15 minutes] (http://www2c.cdc.gov/podcasts/player.asp?f=8642670)</u>
- Vital Signs Motor Vehicle Crash Deaths [PSA 0:60 seconds] (http://www2c.cdc.gov/podcasts/player.asp?f=8642708)
- CDC: Motor Vehicle Safety (http://www.cdc.gov/motorvehiclesafety/)
- CDC: State-based Costs of Motor Vehicle Crash Deaths
 (http://www.cdc.gov/motorvehiclesafety/statecosts/index.html)
- <u>Vital Signs: Motor Vehicle Crash Injuries (http://www.cdc.gov/vitalsigns/crash-injuries/index.html)</u>
- Vital Signs: Child Passenger Safety (http://www.cdc.gov/vitalsigns/childpassengersafety/)
- CDC: Child Passenger Safety (http://www.cdc.gov/Motorvehiclesafety/Child Passenger Safety/index.html)
- CDC: Seat Belts (http://www.cdc.gov/motorvehiclesafety/seatbelts/index.html)
- CDC: Impaired Driving (http://www.cdc.gov/Motorvehiclesafety/Impaired Driving/index.html)
- CDC: Increasing Alcohol Ignition Interlock Use: Successful Practices for States
 (http://www.cdc.gov/motorvehiclesafety/impaired driving/ignition interlock states.html)

- Sobering Facts: Drunk Driving State Fact Sheets
 (http://www.cdc.gov/motorvehiclesafety/impaired driving/states.html)
- CDC: Teen Drivers (http://www.cdc.gov/Motorvehiclesafety/Teen Drivers/index.html)
- CDC: Global Road Safety (http://www.cdc.gov/motorvehiclesafety/global/index.html)
- Prevention Status Reports: Motor Vehicle Injuries (http://www.cdc.gov/psr/national-summary/mvi.html)
- The Community Guide: Motor Vehicle-Related Injury Prevention (http://www.thecommunityguide.org/mvoi/index.html)
- <u>CDC WISQARS (Web-based Injury Statistics Query and Reporting System)</u> (http://www.cdc.gov/injury/wisqars/index.html)
- <u>CDC: Motor Vehicle PICCS (Prioritizing Interventions and Cost Calculator for States)</u> (http://www.cdc.gov/motorvehiclesafety/calculator)

File Formats Help:

How do I view different file formats (PDF, DOC, PPT, MPEG) on this site? (//www.cdc.gov/Other/plugins/)

(//www.cdc.gov/Other/plugins/#pdf)

Page last reviewed: July 6, 2016 Page last updated: July 18, 2016

Content source: Centers for Disease Control and Prevention (http://www.cdc.gov/), National Center for Injury Prevention and

Control (http://www.cdc.gov/injury), Division of Unintentional Injury Prevention Page maintained by: Office of the Associate Director for Communications (OADC)